

vitality impacting the future – today



NASA Johnson Space Center
Astromaterials Research & Exploration Science



THE UNIVERSITY
OF ARIZONA



UNIVERSITY OF
MARYLAND



Calculated Thicknesses of Volcanically Derived Water Ice Deposits at the Lunar Poles

Debra Needham

NASA/Marshall Space Flight Center
and

Matthew Siegler
SMU/PSI

Shuai Li
U. Hawaii at Manoa


David Kring
CLSE/SSERVI – LPI

50th Lunar and Planetary Science Conference
March 22, 2019



Overview

- What we know:
 - Volcanically derived volatiles.
 - Timing of volatile release.
 - Current observations of lunar polar volatiles.
- How volatiles migrated on the Moon.
- Thickness of resulting deposits.
- Implications for the current distribution of lunar volatiles.







CENTER FOR LUNAR SCIENCE AND EXPLORATION

vitality impacting the future – today

Volcanic Volatiles Released from the Moon

- Apollo mare basalt samples: CO, H, OH, H₂O, and S volatiles.
 - e.g., Housley, 1978; Robinson and Taylor, 2014; McCubbin et al., 2010; Shearer et al., 2006
- Volcanic activity peaked 3.8 Ga and 3.5 Ga.
 - 60% of all volcanically derived volatiles were released 3.5 Ga.
 - 20% released 3.8 Ga; remaining 20% released during all other mare eruptions.
- Peak volatile release may have resulted in the formation of a transient lunar atmosphere.
 - Some volatiles lost to space, others settled to the surface as atmosphere dissipated.



Based on data presented in Hiesinger et al., 2011; Whitten et al., 2011


Needham and Kring, 2017, *EPSL*.



CENTER FOR LUNAR SCIENCE AND EXPLORATION

vitaly impacting the future - today

Recent Polar H-Bearing Material Detections from Orbit



Moon Mineralogy Mapper
surface detection of OH/H₂O
at lunar poles (blue/purple);
Pieters et al., 2009


LEND detection of water
equivalent H (1 m depth) via
neutron suppression at lunar
poles; Sanin et al., 2017

OH/H₂O variability by
latitude and day from M³;
Li and Milliken, 2017




Where Lunar Volatiles are Now: North Pole


- Water at surface: Centered about North Pole.
 - Modern accumulations?
- Water at 1 m depth: Offset to 90°E – 180°E .
 - Ancient accumulations? (e.g., Siegler et al., 2016)




Current M³ H₂O Ice
Detection (surface)
Li et al., 2018



LEND Water Equivalent H
(1 m depth)
Sanin et al., 2017



Current Ice Stability Zones
(2.5 m depth)
Siegler et al., 2016




Past Ice Stability Zones
(2.5 m depth)
Siegler et al., 2016




Where Lunar Volatiles are Now: South Pole


- Water at surface: Centered about South Pole.
 - Modern accumulations?
- Water at 1 m depth: Offset to $270^{\circ}\text{E} - 0^{\circ}\text{E}$. (?)
 - Ancient accumulations? (e.g., Siegler et al., 2016)




Current M3 H₂O Ice
Detection (surface)
Li et al., 2018



LEND Water Equivalent H
(1 m depth)
Sanin et al., 2017



Current Ice Stability Zones
(2.5 m depth)
Siegler et al., 2016



Past Ice Stability Zones
(2.5 m depth)
Siegler et al., 2016



Volcanic Volatiles Released from the Moon

- Questions:
 - Where did the volatiles settle on the Moon?
 - How thick would the resulting deposits have been?
- Results have implications for the current distribution of lunar volatiles.






CENTER FOR LUNAR SCIENCE AND EXPLORATION



vitality impacting the future – today

Migration of Lunar Volcanically Derived Volatiles

- In the absence of a lunar atmosphere:
 - Volatiles ‘hop’ based on energy gradient, traveling towards lower energy (to the poles).
 - Assume erupted volatiles migrated to nearest pole – dependent on eruption location.



Releasing volatiles via sputtering and impact vaporization processes; Farrell et al., 2015.



Released volatiles lost to space vs. bounded to the Moon; Farrell et al., 2015.



Migration of Lunar Volcanically Derived Volatiles

- In the presence of a lunar atmosphere:
 - Volatiles entrained in globally distributed atmosphere.
 - Equatorial and mid-latitude volatiles likely to migrate to the poles (e.g., Soto et al., 2018)
 - Assume erupted volatiles deposit evenly at each pole as the atmosphere dissipates – 50% erupted volatiles to each pole.
- Volatiles trapped in areas of stability.





CENTER FOR LUNAR SCIENCE AND EXPLORATION



vitality impacting the future – today

Max Equivalent Thickness of H-Bearing Volcanic Volatile Deposits

Assumptions:


- Volatiles released 3.5 Ga and 3.8 Ga (~80%) split between poles.
- All other volatiles migrated to nearest pole (mostly north pole).
- Assume no H₂O loss (2.4×10^{14} kg) – max deposit thickness.
 - H₂O/OH only; assume H is lost to space
- Know areas of expected volatile preservation (NP/SP):

Region	NP Area (km ²)	SP Area (km ²)	Reference
Current PSRs	12866	16055	Mazarico et al., 2011
Currently Stable 2.5 m	94565	90884	Siegler et al., 2016
Past Stable 2.5 m	86285	82772	Siegler et al., 2016
Observed Surface Water	35	115	Li et al., 2018




Needham and Kring, 2017, *EPSL*.


SP PSRs
Mazarico et
al., 2011




SP Currently
Stable (2.5 m)
Siegler et al.,
2016



SP Current
Surface Ice
Li et al., 2018



SP Past
Stable (2.5 m)
Siegler et al.,
2016







Max Equivalent Thickness of H-Bearing Volcanic Volatile Deposits


- Assumptions:
 - Volatiles released 3.5 Ga and 3.8 Ga (~80%) split between poles.
 - All other volatiles migrated to nearest pole (mostly north pole).
 - Assume no H₂O loss (2.4×10^{14} kg) – max deposit thickness.
 - H₂O/OH only; assume H is lost to space
 - Know areas of expected volatile preservation (NP/SP):

Region	NP Area (km ²)	SP Area (km ²)	Reference
Current PSRs	12866	16055	Mazarico et al., 2011
Currently Stable 2.5 m	94565	90884	Siegler et al., 2016
Past Stable 2.5 m	86285	82772	Siegler et al., 2016




Needham and Kring, 2017, *EPSL*.


SP PSRs
Mazarico et al., 2011




SP Currently
Stable (2.5 m)
Siegler et al.,
2016



SP Current H
Distribution
Sanin et al., 2017



SP Past
Stable (2.5 m)
Siegler et al.,
2016







CENTER FOR LUNAR SCIENCE AND EXPLORATION

vitality impacting the future – today

Max Equivalent Thickness of H-Bearing Volcanic Volatile Deposits


- South Pole Results:

Region	Area (km ²)	Area Reference	Equiv. Thickness (m)
Current PSRs	16055	Mazarico et al., 2011	7.18
Polar Wander Present Stable to 2.5 m	90884	Siegler et al., 2016	1.39
Polar Wander Past Stable to 2.5 m	82772	Siegler et al., 2016	1.27




Needham and Kring, 2017, *EPSL*.


SP PSRs
Mazarico et al., 2011




SP Currently
Stable (2.5 m)
Siegler et al.,
2016



SP Current H
Distribution
Sanin et al., 2017



SP Past
Stable (2.5 m)
Siegler et al.,
2016







CENTER FOR LUNAR SCIENCE AND EXPLORATION

vitality impacting the future – today

Maximum Thickness of H-Bearing Volcanic Volatile Deposits


- North Pole Results:

Region	Area (km ²)	Area Reference	Equiv. Thickness (m)
Current PSRs	12866	Mazarico et al., 2011	9.70
Polar Wander Present Stable to 2.5 m	94565	Siegler et al., 2016	1.45
Polar Wander Past Stable to 2.5 m	86285	Siegler et al., 2016	1.32




Needham and Kring, 2017, *EPSL*.


NP PSRs
Mazarico et al., 2011




NP Currently
Stable (2.5 m)
Siegler et al.,
2016



NP Current H
Distribution
Sanin et al., 2017



NP Past
Stable (2.5 m)
Siegler et al.,
2016







CENTER FOR LUNAR SCIENCE AND EXPLORATION

vitality impacting the future – today


Implications for Distribution of Polar Lunar Volatiles

- Ice ~1.5 m thick deposited in stable regions at each lunar pole.
 - Subsequently covered by ejecta, vaporized, and gardened by subsequent impacts.
 - Expected to have 6-10 m ice-bearing regolith above thinner subsurface ice horizon.

(Fa and Jin, 2010; Kobayashi et al., 2010)



Releasing volatiles via sputtering and impact vaporization processes; Farrell et al., 2015.



Released volatiles lost to space vs. bounded to the Moon; Farrell et al., 2015.




CENTER FOR LUNAR SCIENCE AND EXPLORATION

vitality impacting the future – today

Implications for a Mission Prospecting for Lunar Volatiles


- The source of volatiles can affect the composition of these volatile deposits.
 - H, O isotopes
 - Alteration minerals like hematite! (Li et al., this meeting)




Solar wind-delivered hydrogen trapped in the Moon's PSRs.



Water delivered by asteroid and comet impacts on the Moon.




Water-building components erupted during volcanic eruptions.



D/H isotopes across the Solar System, from Saal et al., 2013 and references therein.

Oxygen isotopes across the Solar System from Righter and O'Brien, 2007 and refs. therein.






CENTER FOR LUNAR SCIENCE AND EXPLORATION

vitality impacting the future – today

Implications for a Mission Prospecting for Lunar Volatiles


- The source of volatiles can affect the composition of these volatile deposits.
 - H, O isotopes
 - Alteration minerals like hematite! (Li et al., this meeting)






Solar wind-delivered hydrogen trapped in the Moon's PSRs.



Water delivered by asteroid and comet impacts on the Moon.



Water delivered by asteroid and comet impacts on the Moon.





CENTER FOR LUNAR SCIENCE AND EXPLORATION

vitality impacting the future – today

